

The EIC, Saturation, and the CGC

Thomas Ullrich (BNL)
RIKEN Glasma Workshop, BNL
Wednesday, May 12, 2010

The EIC, Saturation, and the CGC

Thomas Ullrich (BNL)
RIKEN Glasma Workshop, BNL
Wednesday, May 12, 2010



EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
 - ▶ Study the Physics of Strong Color Fields
 - ◉ Establish (or not) the existence of the saturation regime
 - ◉ Explore non-linear QCD
 - ◉ Measure momentum & space-time distributions of glue
 - ▶ Study the nature of color singlet excitations (Pomerons)
 - ▶ Test and study the limits of universality (eA vs. pA)

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons
 - ▶ What is the nature of the spin of the proton?
 - ⊙ $\Delta G(Q^2, x)$, polarization of the sea quarks
 - ⊙ Transverse spin and momentum measurements and correlations
 - ▶ What is the Three-Dimensional Spatial Landscape of Nucleons?
 - ⊙ Transverse imaging of quarks and gluons in nucleons
 - ⊙ Transverse momentum dependent measurements and correlations

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons
- What governs the transition of quarks and gluons into pions and nucleons?
 - ▶ How do fast probes interact with the gluonic medium?
 - ⦿ Energy loss of quarks and gluons
 - ▶ Mechanism of fragmentation?

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons
- What governs the transition of quarks and gluons into pions and nucleons?

EIC Science Case

How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons
- What governs the transition of quarks and gluons into pions and nucleons?
- Electroweak Physics (studies underway)
 - ▶ Parity Violating deep inelastic scattering (PVDIS)
 - ◉ Quark helicity distributions
 - ◉ Isovector EMC effect
 - ◉ Potential ultraprecise weak mixing angle measurements
 - ▶ Lepton Flavor and Number Violation
 - ◉ Electron-tau lepton conversion

EIC Science Case


How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

- What is the nature and role of gluons and their self-interactions (eA, ep)
- What is the internal landscape of the nucleons
- What governs the transition of quarks and gluons into pions and nucleons?
- Electroweak Physics (studies underway)

EIC Science Case


How do we understand the visible matter in our universe in terms of the fundamental quarks and gluons of QCD?

The machine presents a unique opportunity for fundamental physics:



Why Electrons ?


Hadron-Hadron



- Test QCD
- Probe/Target interaction directly via gluons
- Lacks the direct access to parton kinematics
- Probe has complex structure


Why Electrons ?

Hadron-Hadron




- Test QCD
- Probe/Target interaction directly via gluons
- Lacks the direct access to parton kinematics
- Probe has complex structure

Electron-Hadron (DIS)




- Explore QCD & Hadron Structure
- Indirect access to glue
- High precision & access to partonic kinematics
- Probes partons w/o disturbing them or interfering with their dynamics

Why Electrons ?



Electron-Hadron (DIS)




- Explore QCD & Hadron Structure
- Indirect access to glue
- High precision & access to partonic kinematics
- Probes partons w/o disturbing them or interfering with their dynamics

Both are complementary but for precision \Rightarrow ep, eA


Gluon Distributions and Saturation

How to probe saturation? $G(x, Q^2)$ is not an observable!



Gluon Distributions and Saturation

How to probe saturation? $G(x, Q^2)$ is not an observable!





Comparison (to constrain/reject models) requires

- ▶ “lever arm” in x , Q^2 , A , ...
- ▶ complementary measurements (incl., semi-incl., excl., jets, DIS & diffractive, varying probes, ...)

EIC - Reaching the Saturation Regime

Saturation:

- Au: Strong hints from RHIC at $x \sim 10^{-3}$
- p: No (?) hints at Hera up to $x=6.32 \cdot 10^{-5}$, $Q^2 = 1-5 \text{ GeV}^2$




EIC - Reaching the Saturation Regime

Saturation:

- Au: Strong hints from RHIC at $x \sim 10^{-3}$
- p: No (?) hints at Hera up to $x=6.32 \cdot 10^{-5}$, $Q^2 = 1-5 \text{ GeV}^2$

Nuclear Enhancement:

$$(Q_s^A)^2 \approx c Q_0^2 \left(\frac{A}{x} \right)^{1/3}$$




EIC - Reaching the Saturation Regime

Saturation:

- Au: Strong hints from RHIC at $x \sim 10^{-3}$
- p: No (?) hints at Hera up to $x=6.32 \cdot 10^{-5}$, $Q^2 = 1-5 \text{ GeV}^2$

Nuclear Enhancement:

$$(Q_s^A)^2 \approx c Q_0^2 \left(\frac{A}{x} \right)^{1/3}$$




EIC - Reaching the Saturation Regime

Saturation:

- Au: Strong hints from RHIC at $x \sim 10^{-3}$
- p: No (?) hints at Hera up to $x=6.32 \cdot 10^{-5}$, $Q^2 = 1-5 \text{ GeV}^2$

Nuclear Enhancement:

$$(Q_s^A)^2 \approx c Q_0^2 \left(\frac{A}{x} \right)^{1/3}$$



EIC - Reaching the Saturation Regime


Saturation:

- Au: Strong hints from RHIC at $x \sim 10^{-3}$
- p: No (?) hints at Hera up to $x=6.32 \cdot 10^{-5}$, $Q^2 = 1-5 \text{ GeV}^2$


Nuclear Enhancement:

$$(Q_s^A)^2 \approx c Q_0^2 \left(\frac{A}{x} \right)^{1/3}$$

Finding RHIC and Hera
& Q_s scalings consistent



EIC - Energy Requirements




Coverage

- Need lever arm in Q^2 at fixed x to constrain models
- Need $Q > Q_s$ to study onset of saturation
- $Q^2 = 4EE'\sin^2\theta/2$
 - ▶ hard to get much lower than $0.5\text{-}1 \text{ GeV}^2$

- ep: $\sim 1 \text{ TeV}$ is needed
- eA: $\sqrt{s} = 50 \text{ GeV}$ is marginal, around $\sqrt{s} = 100 \text{ GeV}$ desirable

EIC - Energy Requirements




Coverage

- Need lever arm in Q^2 at fixed x to constrain models
- Need $Q > Q_s$ to study onset of saturation
- $Q^2 = 4EE'\sin^2\theta/2$
 - ▶ hard to get much lower than 0.5-1 GeV²

- ep: ~ 1 TeV is needed
- eA: $\sqrt{s} = 50$ GeV is marginal, around $\sqrt{s} = 100$ GeV desirable

EIC - Energy Requirements



Coverage

- Need lever arm in Q^2 at fixed x to constrain models
- Need $Q > Q_s$ to study onset of saturation
- $Q^2 = 4EE'\sin^2\theta/2$
 - ▶ hard to get much lower than 0.5-1 GeV^2


Current best estimate to study saturation:

Electron energy $\sim 20 \text{ GeV}$ x Ion Energy $\sim 100 \text{ GeV}$

EIC - What Luminosity is Needed?

Hera suffered from low luminosity: $L = 1.6\text{-}3.8 \cdot 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$

EIC-Example F_L : *inclusive* measurements at different \sqrt{s} , assume 1% energy-to-energy normalization



$\int \mathcal{L} dt = 4/A \text{ fb}^{-1}$ (10+100) GeV &
 $4/A \text{ fb}^{-1}$ (10+50) GeV &
 $2/A \text{ fb}^{-1}$ (5+50) GeV


All together 10 weeks at
 $L \sim 4 \cdot 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ and 50% duty
cycle
(Note: 100x Hera L)

Conclusion from this
study:


- Dominated by **sys. uncertainties**
- More statistics does not provide higher accuracy

EIC - What Luminosity is Needed?

Key measurement: exclusive diffractive vector meson production: $e A \rightarrow e' A' V$



Sensitive means to probe $G(x, Q^2)$
and saturation: $\sigma \propto G(x, Q^2)^2$



$\sigma(x_{IP}, x, Q^2, t)$ double/triple differential & small σ
 \Rightarrow **high L required** (statistical \gg systematic errors),
 (DVCS even more L hungry \rightarrow GPDs)

EIC - Requirements

Requirements depend on specific measurements

| Measurement | | \sqrt{s} | L (cm ⁻² s ⁻¹) |
|---|---|---|---|
| Saturation (eA) | Inclusive F2, FL | High $\sqrt{s} > 50$ GeV preferred $\sqrt{s} \sim 100$ GeV | moderate $\sim 10^{32}$ |
| | Diffraction VM & n-differential | | high $\geq 10^{33}$ |
| 3D Imaging (ep and eA) | GPD, TMD n-differential | Moderate $\sqrt{s} > 50$ GeV | high $\sim 10^{34}$ |
| Electroweak (ep) | PVDIS, W production, γ -Z interference, Lepton number violation | Moderate to high | very high $\geq 10^{34}$ |
| Spin structure of the nucleon (ep) | Inclusive: structure functions, g_1 , ... Semi-inclusive (tag quark) | High $\sqrt{s} > 50$ GeV preferred $\sqrt{s} \sim 100$ GeV | moderate-high $\sim 10^{33}$ |
| Fragmentation, Energy loss (ep, eA) | Spectra: ν , p_T , Q^2 flavor dependence (c, b) | Low - High $\sqrt{s} > 50$ GeV (heavy flavor) | moderate $\sim 10^{32}$ |

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow staged approach

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow staged approach

- QCD Community: JLAB & RHIC
 - ▶ BNL: EIC Task Force, eRHIC-CAD
 - ▶ JLAB: User Group, Accelerator Department

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow staged approach

- **QCD Community**: JLAB & RHIC
 - ▶ BNL: EIC Task Force, eRHIC-CAD
 - ▶ JLAB: User Group, Accelerator Department
- **EIC Collaboration** (<http://web.mit.edu/eicc/>)
 - ▶ Collaboration Meetings, Working Groups
 - ▶ Steering Committee

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow staged approach

- **QCD Community**: JLAB & RHIC
 - ▶ BNL: EIC Task Force, eRHIC-CAD
 - ▶ JLAB: User Group, Accelerator Department
- **EIC Collaboration** (<http://web.mit.edu/eicc/>)
 - ▶ Collaboration Meetings, Working Groups
 - ▶ Steering Committee
- **International Advisory Committee**
 - ▶ Members from broad community, chair: W. Henning

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow staged approach


- **QCD Community**: JLAB & RHIC
 - ▶ BNL: EIC Task Force, eRHIC-CAD
 - ▶ JLAB: User Group, Accelerator Department
- **EIC Collaboration** (<http://web.mit.edu/eicc/>)
 - ▶ Collaboration Meetings, Working Groups
 - ▶ Steering Committee
- **International Advisory Committee**
 - ▶ Members from broad community, chair: W. Henning
- **EIC Concepts**
 - ▶ eRHIC (RHIC/BNL)
 - ▶ ELIC (CEBAF/JLAB)

EIC - Realization

High \sqrt{s} , high L machine: costs too high to build in one step \Rightarrow **staged approach**

- **QCD Community**: JLAB & RHIC
 - ▶ BNL: EIC Task Force, eRHIC-CAD
 - ▶ JLAB: User Group, Accelerator Department
- **EIC Collaboration** (<http://web.mit.edu/eicc/>)
 - ▶ Collaboration Meetings, Working Groups
 - ▶ Steering Committee
- **International Advisory Committee**
 - ▶ Members from broad community, chair: W. Henning
- **EIC Concepts**
 - ▶ eRHIC (RHIC/BNL)
 - ▶ ELIC (CEBAF/JLAB)
- **Timeline**: ~2020 (vital: NSAC Long Range Plan 2012/13)

EIC Concept: CEBAF → ELIC @ JLAB



EIC Concept: CEBAF → ELIC @ JLAB

Use CEBAF “as-is” after 12-GeV Upgrade

MEIC:


- 3-11 GeV e
- 60 GeV p ($\sqrt{s} = 51$ GeV)
- 24 GeV/n ion ($\sqrt{s} = 33$ GeV)
- $L \sim 1 \cdot 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ @ 5+60 GeV
- $L \sim 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ otherwise
- Up to heavy ion $A = 208$

ELIC: upgrades


- 250 GeV p ($\sqrt{s} = 105$ GeV)
- 100 GeV ion ($\sqrt{s} = 66$ GeV)
- L up to $10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

ELIC: upgrades

- Increase E_e



EIC Concept: RHIC \rightarrow eRHIC @ BNL




EIC Concept: RHIC \rightarrow eRHIC @ BNL

ERL: Energy Recovery Linac



1 - 5 GeV per pass

4 (6) passes






EIC Concept: RHIC \rightarrow eRHIC @ BNL

ERL: Energy Recovery Linac
1 - 5 GeV per pass
4 (6) passes



EIC Concept: RHIC \rightarrow eRHIC @ BNL

ERL: Energy Recovery Linac
1 - 5 GeV per pass
4 (6) passes



EIC Concept: RHIC \rightarrow eRHIC @ BNL

eRHIC (stage 1):


- 4 GeV e \times 250 GeV p (\sqrt{s} = 63 GeV)
- 4 GeV e $^-$ \times 100 GeV/n Au (\sqrt{s} = 40 GeV)
- $L \sim 10^{32}$ - 10^{33} cm $^{-2}$ s $^{-1}$

eRHIC (stage 2):

- 20 GeV e \times 325 GeV p (\sqrt{s} = 160 GeV)
- 20 GeV e \times 130 GeV/n Au (\sqrt{s} = 102 GeV)
- $L \sim 1.4 \cdot 10^{34}$ cm $^{-2}$ s $^{-1}$

eRHIC upgrades

- Higher luminosity
- Higher hadron energy




eRHIC: Detector \Leftrightarrow IR Design

Detector:

- Hermeticity (diffraction \Leftrightarrow rapidity gap)
- Extreme forward angle \Leftrightarrow low-x physics
- $PID < 4 \text{ GeV}/c$ in central detector
- PID up to 100 GeV for forward rapidities
- High precision for scattered electron that defines event kinematics (Q^2 , x , ...) \Rightarrow low material

Detector concept (schematic):




eRHIC: Detector \Leftrightarrow IR Design

IR:

- minimize synchrotron radiation (P and λ)
- minimize $\beta^* \Rightarrow L$
- First quads: maximize aperture for particles at extreme forward angle \Rightarrow detect n from breakup (diffraction)
- Design depends on presence/absence of pp/pA/AA ops

High Luminosity IR ($\beta^ = 5$ cm)*



Dejan Trbojevic

Latest Design:



- ep(A) crossing angle of 10 mrad
- straight e beamline
 \Rightarrow no synchrotron radiation
- good separation of e/p beamlines
 - real estate for additional detectors
- need to fine tune for optimal instrumentation
- aperture of 1st quad: ± 5 mrad
 - important for breakup-neutron detection

Heavy Ion Experiments and ep/eA?

eRHIC & staged eRHIC:

- Electron beam around RHIC tunnel
- Requires scheme for getting electron beamline **through** or **around** PHENIX and STAR
- Can STAR/PHENIX make use of the e-p and e-A collisions?
 - ▶ **RHIC decadal planing:** experiments are looking into it

Example: STAR



EIC Outlook


- Increasing efforts at BNL & JLAB
- Concepts of eRHIC and ELIC are taking shape
 - ▶ Substantial progress in machine & IR design
- Staged approach most promising path
 - ▶ Much can be done already at lower energy (e.g. F_L)
 - ▶ **Saturation** physics will require **full ELIC/eRHIC**
- At minimum one large multi-purpose detector
- ePHENIX/eSTAR under evaluation

Key Events:

- INT Workshop: Gluons and the Quark Sea at High Energies: Sep-Nov, 2010 \Rightarrow Science Case for EIC
- LRP 2012: Town meetings and LRP

Additional Slides

Past and Future Electron-Hadron Collider



Very long term future

- LHeC: add linac/ring to existing LHC
 - ▶ 70 GeV e-beam on 2.8 TeV Pb (7 TeV p), $\sqrt{s_{eN}} = 885$ GeV
- ENC@FAIR: add e-ring to planned HESR p ring
 - ▶ 3.3 GeV e-beam on 15 GeV p, $\sqrt{s} = 14$ GeV

Measurements & Techniques

- Gluon Distribution $G(x, Q^2)$

- ▶ Scaling violation in F_2 : $\delta F_2 / \delta \ln Q^2$
- ▶ $F_L \sim xG(x, Q^2)$
- ▶ 2+1 jet rates
- ▶ Diffractive vector meson production ($[xG(x, Q^2)]^2$)

- Space-Time Distribution

- ▶ Exclusive diffractive VM production ($J/\psi, \phi, \rho$)
- ▶ Deep Virtual Compton Scattering (nGPDs)
- ▶ Structure functions for various mass numbers A and its impact parameter dependence